



DECLARATION

I, Emiko Oku of HIROTA & ASSOCIATES, residing at Wakabayashi Bldg. 3F, 8-5, Akasaka 2-chome, Minato-ku, Tokyo 107-0052, Japan, do hereby certify that I am conversant with the English and Japanese languages and am a competent translator thereof, and I further certify that to the best of my knowledge and belief the following is a true and correct translation made by me of the document in the Japanese language filed for a patent application in Japan under No. 11-368991 on December 27, 1999 in the name of JAPAN SCIENCE AND TECHNOLOGY CORPORATION at Tokyo, Japan, entitled: "HIGH-AFFINITY CHOLINE TRANSPORTER".

Signed this 29 day of August, 2006

A handwritten signature in cursive script, appearing to read "Emiko Oku", written over a horizontal line.

Emiko Oku



PATENT OFFICE
JAPANESE GOVERNMENT

This is to certify that the annexed is a true copy of the following application as filed with this Office.

Date of Application: December 27, 1999

Application Number: Patent Application No.11-368991

Applicant(s): JAPAN SCIENCE AND TECHNOLOGY
CORPORATION

Commissioner, Kozo Oikawa (Sealed)
Patent Office

【Name of Document】	Patent Application
【Case Number】	A011P15
【Filing Date】	December 27, 1999
【Addressee】	Commissioner of Patent Office
【International Classification】	C12N 15/00
【Inventor】	
【Address or Residence】	Room 411, 3-1, Numama 2-chome, Zushi-shi, Kanagawa
【Name】	Tatsuya HAGA
【Inventor】	
【Address or Residence】	Room 101, Rizu Hibarigaoka, 1-6, Hibarigaokakita 4-chome, Hoya-shi, Tokyo
【Name】	Takashi OKUDA
【Applicant for Patent】	
【Code Number】	396020800
【Name or Corporation Name】	JAPAN SCIENCE AND TECHNOLOGY CORPORATION
【Representative】	Moritaka NAKAMURA
【Agent】	
【Code Number】	100107984
【Patent Attorney】	
【Name or Corporation Name】	Masanori HIROTA
【Priority Claim Based on Previous Application】	
【Application Number】	Application No. 11-240642
【Filing Date】	August 27, 1999
【Indication of Fee】	
【Deposit Account Number】	044347
【Amount of Payment】	21,000 yen
【List of Filed Documents】	

[Name of Document]	Specification 1
[Name of Document]	Drawing 1
[Name of Document]	Abstract 1
[Confirmation of Proof]	Proof Necessary

[Name of Document] SPECIFICATION

[Title of the Invention] HIGH-AFFINITY CHOLINE TRANSPORTER

[Scope of Claims]

[Claim 1] A gene which encodes a protein having high-affinity choline transporter activity.

[Claim 2] A gene which encodes a protein (a) or (b) described below;

(a) a protein comprising an amino acid sequence represented by Seq. ID No. 2,

(b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.2, and having high-affinity choline transporter activity.

[Claim 3] DNA containing a base sequence represented by Seq. ID No. 1 or its complementary sequence and a part or a whole of these sequences.

[Claim 4] DNA derived from a nematode which hybridizes with DNA comprising a gene according to claim 3 under a stringent condition, and encodes a protein having high-affinity choline transporter activity.

[Claim 5] A gene which encodes a protein (a) or (b) described below;

(a) a protein comprising an amino acid sequence represented by Seq. ID No. 4,

(b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.4, and having high-affinity choline transporter activity.

[Claim 6] DNA containing a base sequence represented by Seq. ID No. 3 or its complementary sequence and a part or a whole

of these sequences.

【Claim 7】 DNA derived from a rat which hybridizes with DNA comprising a gene according to claim 6 under a stringent condition, and encodes a protein having high-affinity choline transporter activity.

【Claim 8】 A gene which encodes a protein (a) or (b) described below;

(a) a protein comprising an amino acid sequence represented by Seq. ID No. 6,

(b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.6, and having high-affinity choline transporter activity.

【Claim 9】 DNA containing a base sequence represented by Seq. ID No. 5 or its complementary sequence and a part or a whole of these sequences.

【Claim 10】 DNA derived from a human which hybridizes with DNA comprising a gene according to claim 9 under a stringent condition, and encodes a protein having high-affinity choline transporter activity.

【Claim 11】 A gene which encodes a protein (a) or (b) described below;

(a) a protein comprising an amino acid sequence represented by Seq. ID No. 8,

(b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.8, and having high-affinity choline transporter activity.

【Claim 12】 DNA containing a base sequence represented by Seq. ID No. 7 or its complementary sequence and a part or a whole

of these sequences.

[Claim 13] DNA derived from a mouse which hybridizes with DNA comprising a gene according to claim 12 under a stringent condition, and encodes a protein having high-affinity choline transporter activity.

[Claim 14] A protein having high-affinity choline transporter activity.

[Claim 15] A protein comprising an amino acid sequence represented by Seq. ID No. 2.

[Claim 16] A protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 2, and having nematode high-affinity choline transporter activity.

[Claim 17] A protein comprising an amino acid sequence represented by Seq. ID No. 4.

[Claim 18] A protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 4, and having rat high-affinity choline transporter activity.

[Claim 19] A protein comprising an amino acid sequence represented by Seq. ID No. 6.

[Claim 20] A protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 6, and having human high-affinity choline transporter activity.

[Claim 21] A protein comprising an amino acid sequence represented by Seq. ID No. 8.

[Claim 22] A protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 8, and having

mouse high-affinity choline transporter activity.

[Claim 23] A fusion protein being constructed by expressing a cDNA encoding fusion proteins of a protein having high-affinity choline transporter activity and a marker protein and/or a peptide tag.

[Claim 24] The fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16.

[Claim 25] The fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18.

[Claim 26] The fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20.

[Claim 27] The fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22.

[Claim 28] An antibody which specifically binds to a protein having high-affinity choline transporter activity.

[Claim 29] The antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16.

[Claim 30] The antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to

claim 17 or 18.

[Claim 31] The antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20.

[Claim 32] The antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22.

[Claim 33] The antibody according to any one of claims 28 to 32, wherein the antibody is a monoclonal antibody.

[Claim 34] A host cell containing an expression system which can express a protein having high-affinity choline transporter activity.

[Claim 35] The host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16.

[Claim 36] The host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18.

[Claim 37] The host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20.

[Claim 38] The host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22.

[Claim 39] A non-human animal whose function of a gene which encodes a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome.

[Claim 40] The non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16.

[Claim 41] The non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18.

[Claim 42] The non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20.

[Claim 43] The non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22.

[Claim 44] The non-human animal according to any one of claims 39 to 43, wherein the non-human animal is a mouse or a rat.

[Claim 45] A preparing method of a cell having high-affinity choline transporter activity characterized in introducing the gene or the DNA according to any one of claims 8 to 10 into a cell whose function of a gene which encodes a protein having high-affinity choline transporter activity is deficient on its chromosome.

[Claim 46] The preparing method of a cell having high-affinity choline transporter activity according to claim 45, wherein the cell having high-affinity choline transporter activity is

integrated with the gene or the DNA according to any one of claims 8 to 10 in its chromosome, and stably shows high-affinity choline transporter activity.

[Claim 47] A cell having high-affinity choline transporter activity being obtainable by the preparing method of a cell having high-affinity choline transporter activity according to claim 45 or 46.

[Claim 48] A screening method of a promoter or a suppressor of high-affinity choline transporter activity characterized in measuring/evaluating high-affinity choline transporter activity of the protein having high-affinity choline transporter activity according to any one of claims 14 to 22 in the presence of a subject material.

[Claim 49] A screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in comprising the steps of: a cell membrane or a cell which expresses a protein having high-affinity choline transporter activity is cultivated in vitro in the presence of a subject material; the activity and/or the expression amount of a protein having high-affinity choline transporter activity in the cell membrane or the cell is measured/evaluated.

[Claim 50] The screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to claim 49, wherein the cell membrane or the cell which expresses a protein having high-affinity choline transporter activity is the host cell containing an expression system which can express a protein having high-affinity choline transporter activity according to any one of claims 34 to 38, or is the cell having high-affinity

choline transporter activity according to claim 47.

[Claim 51] The screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to any one of claims 48 to 50, wherein the protein having high-affinity choline transporter activity is a recombinant protein.

[Claim 52] A screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in comprising the steps of: a cell obtained from the non-human animal according to any one of claims 39 to 44 is cultivated in vitro in the presence of a subject material; the activity and/or the expression amount of a protein having high-affinity choline transporter activity in the cell is measured/evaluated.

[Claim 53] A screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity.

[Claim 54] A screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome, and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity.

[Claim 55] A screening method of a promoter or a suppressor

of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome, and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity in comparison with the case using wild-type non-human animal.

[Claim 56] The screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to any one of claims 52 to 55, wherein the non-human animal is a mouse or a rat.

[Claim 57] A material which promotes activity or expression of a protein having high-affinity choline transporter activity being obtainable by the screening method according to any one of claims 48 to 56.

[Claim 58] A material which suppresses activity or expression of a protein having high-affinity choline transporter activity being obtainable by the screening method according to any one of claims 48 to 56.

[Claim 59] A medical constituent characterized in being used for a medical treatment for a patient who needs elevation of the activity or enhancement of the expression of a high-affinity choline transporter, and containing the protein according to any one of claims 14 to 22, and/or the material which promotes activity or expression of a protein having high-affinity choline transporter activity according to claim 57 as an active component.

[Claim 60] A medical constituent characterized in being used for medical treatment for a patient who needs suppression of the activity or the expression of a high-affinity choline transporter, and containing the protein according to any one of claims 14 to 22, and/or the material which suppresses the activity or the expression of a protein having high-affinity choline transporter activity according to claim 58 as an active component.

[Claim 61] A diagnostic method for diseases relating to the expression or the activity of a high-affinity choline transporter characterized in comparing a DNA sequence encoding a high-affinity choline transporter in a sample to a DNA sequence encoding the protein according to claim 19 or 20.

[Claim 62] A diagnostic probe for Alzheimer's disease comprising a whole or a part of an antisense strand of DNA or RNA encoding the protein according to claim 19 or 20.

[Claim 63] A diagnostic drug for Alzheimer's disease characterized in containing the diagnostic probe according to claim 62 and/or the antibody according to any one of claims 28 to 33.

[Detailed Description of the Invention]

[0001]

[Technical Field to Which the Invention Pertains]

This invention relates to a protein having high-affinity choline transporter activity, a gene encoding said protein and the use of the same.

[0002]

[Prior Art]

The autonomic nervous system which spreads to organs throughout a body and regulates the most basic functions of living

organism including energy metabolism, circulation, respiration and reproduction along with endocrine system, is classified into the sympathetic and parasympathetic nervous systems. All autonomic nerve fibers excluding postganglionic fibers of the sympathetic nerve, motor nerve fiber, and sudoriferous gland/blood vessel dilative fiber in the sympathetic nerve are cholinergic, and acetylcholine is vital for the function of the autonomic nerve and the motor nerve. It has been known that the cholinergic neuron, being observed also in the brain, is important for recognizing function of the brain and that it degenerates after the onset of Alzheimer's disease. In the cholinergic neuron, because of lack of biosynthetic ability for choline, choline, an acetylcholine decomposition product, is taken up into a cell by a high-affinity choline transporter at the presynaptic terminals to be reused for synthesizing acetylcholine. The high-affinity choline uptake is a rate-limiting step for acetylcholine synthesis and is presumed to regulate the efficiency of synaptic transmission (J. Neurochem. 18, 781-798, 1971, Science 178, 626-628, 1972, Biochem. Biophys. Acta 291, 564-575, 1973, Mol. Pharmacol. 9, 630-639, 1973, J. Pharmacol. Exp. Ther. 192, 86-94, 1975, J. Neurochem. 30, 15-21, 1978, J. Neurochem. 44, 11-24, 1985, J. Neurochem. 60, 1191-1201, 1993, J. Neurochem. 20, 581-593, 1973, Eur. J. Pharmacol. 102, 369-370, 1984). To date, most of cDNAs of transporters for major neurotransmitters have been isolated, however, a cDNA of the high-affinity choline transporter, which is physiologically important, has not been identified.

[0003]

[An Object to be Attained]

So far, the existence of a protein being localized in the

cholinergic neuron and having a function of taking up choline, a precursor of acetylcholine, into a cell has been expected, but molecular properties of said protein, a high-affinity choline transporter, have been unknown. An object of the present invention is to provide a physiologically important protein having the high-affinity choline transporter activity, a gene which encodes the protein, and a screening method of a high-affinity choline transporter activity promoter using the protein, the gene and the like.

[0004]

[Means to Attain the Object]

The inventors have conducted intensive study to attain the above-mentioned object: with information of genomic project (Science 282, 2012-2018, 1998), Na^+ -dependent transporter cDNAs being expected from the genomic sequence of a nematode (*C. elegans*) were cloned one by one, and the high-affinity choline uptake activity of each cDNA was examined in the oocyte expression system of *Xenopus*, and the cDNA of nematode high-affinity choline transporter (cho-1) was identified on the basis of the above examination, then homologous molecules (CHT1) were cloned from rat spinal cord by using the homology of a base sequence to the cDNA as an index. This CHT1 had no homology to neurotransmitter transporters (J. Neurochem. 71, 1785-1803, 1998), but had 20 to 25% homology to molecules which belong to Na^+ -dependent glucose transporter family (Nature 330, 379-381, 1987).

[0005]

Northern blot analysis revealed that transcripts of CHT1 were confirmed only in spinal cord, basal forebrain, corpus striatum and brain stem, and CHT1 seemed to be expressed in cholinergic neurons. Accordingly, CHT1 was expressed in

oocytes of *Xenopus*. As a result, choline uptake activity that is Na⁺-dependent and completely inhibited by hemicholinium-3 was observed. These results indicate that CHT1 has high-affinity choline transporter activity. Further, the inventors have cloned choline transporter cDNAs derived from a human and from a mouse, and determined their base sequences, and have confirmed that their expression products have high-affinity choline uptake activity. The present invention has thus completed.

[0006]

The present invention relates to a gene which encodes a protein having high-affinity choline transporter activity (claim 1), a gene which encodes a protein (a) or (b) described below; (a) a protein comprising an amino acid sequence represented by Seq. ID No. 2, (b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.2, and having high-affinity choline transporter activity (claim 2), DNA containing a base sequence represented by Seq. ID No. 1 or its complementary sequence and a part or a whole of these sequences (claim 3), DNA derived from a nematode which hybridizes with DNA comprising a gene according to claim 3 under a stringent condition, and encodes a protein having high-affinity choline transporter activity (claim 4), a gene which encodes a protein (a) or (b) described below; (a) a protein comprising an amino acid sequence represented by Seq. ID No. 4, (b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No.4, and having high-affinity choline transporter activity (claim 5), DNA

containing a base sequence represented by Seq. ID No. 3 or its complementary sequence and a part or a whole of these sequences (claim 6), DNA derived from a rat which hybridizes with DNA comprising a gene according to claim 6 under a stringent condition, and encodes a protein having high-affinity choline transporter activity (claim 7), a gene which encodes a protein (a) or (b) described below; (a) a protein comprising an amino acid sequence represented by Seq. ID No. 6, (b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 6, and having high-affinity choline transporter activity (claim 8), DNA containing a base sequence represented by Seq. ID No. 5 or its complementary sequence and a part or a whole of these sequences (claim 9), DNA derived from a human which hybridizes with DNA comprising a gene according to claim 9 under a stringent condition, and encodes a protein having high-affinity choline transporter activity (claim 10), a gene which encodes a protein (a) or (b) described below; (a) a protein comprising an amino acid sequence represented by Seq. ID No. 8, (b) a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 8, and having high-affinity choline transporter activity (claim 11), DNA containing a base sequence represented by Seq. ID No. 7 or its complementary sequence and a part or a whole of these sequences (claim 12), and DNA derived from a mouse which hybridizes with DNA comprising a gene according to claim 12 under a stringent condition, and encodes a protein having high-affinity choline transporter activity (claim 13).

[0007]

The present invention also relates to a protein having high-affinity choline transporter activity (claim 14), a protein comprising an amino acid sequence represented by Seq. ID No. 2 (claim 15), a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 2, and having nematode high-affinity choline transporter activity (claim 16), a protein comprising an amino acid sequence represented by Seq. ID No. 4 (claim 17), a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 4, and having rat high-affinity choline transporter activity (claim 18), a protein comprising an amino acid sequence represented by Seq. ID No. 6 (claim 19), a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 6, and having human high-affinity choline transporter activity (claim 20), a protein comprising an amino acid sequence represented by Seq. ID No. 8 (claim 21), and a protein comprising an amino acid sequence where one or a few amino acids are deficient, substituted or added in the amino acid sequence represented by Seq. ID No. 8, and having mouse high-affinity choline transporter activity (claim 22).

[0008]

The present invention further relates to a fusion protein being constructed by expressing a cDNA encoding fusion proteins of a protein having high-affinity choline transporter activity and a marker protein and/or a peptide tag (claim 23), the fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has nematode

high-affinity choline transporter activity according to claim 15 or 16 (claim 24), the fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18 (claim 25), the fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20 (claim 26), and the fusion protein according to claim 23, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22 (claim 27).

[0009]

The present invention still further relates to an antibody which specifically binds to a protein having high-affinity choline transporter activity (claim 28), the antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16 (claim 29), the antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18 (claim 30), the antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20 (claim 31), the antibody according to claim 28, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22 (claim 32), and the antibody according to any one of claims 28 to 32, wherein the antibody is a monoclonal

antibody (claim 33).

[0010]

The present invention also relates to a host cell containing an expression system which can express a protein having high-affinity choline transporter activity (claim 34), the host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16 (claim 35), the host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18 (claim 36), the host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20 (claim 37), and the host cell according to claim 34, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22 (claim 38).

[0011]

The present invention further relates to a non-human animal in which function of a gene which encodes a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome (claim 39), the non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has nematode high-affinity choline transporter activity according to claim 15 or 16 (claim 40), the non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has rat high-affinity choline transporter activity according to claim 17 or 18 (claim 41), the non-human animal according to claim

39, wherein the protein having high-affinity choline transporter activity has human high-affinity choline transporter activity according to claim 19 or 20 (claim 42), the non-human animal according to claim 39, wherein the protein having high-affinity choline transporter activity has mouse high-affinity choline transporter activity according to claim 21 or 22 (claim 43), and the non-human animal according to any one of claims 39 to 43, wherein the non-human animal is a mouse or a rat (claim 44).

[0012]

The present invention still further relates to a preparing method of a cell having high-affinity choline transporter activity characterized in introducing the gene or the DNA according to any one of claims 8 to 10 into a cell whose function of a gene which encodes a protein having high-affinity choline transporter activity is deficient on its chromosome (claim 45), the preparing method of a cell having high-affinity choline transporter activity according to claim 45, wherein the cell having high-affinity choline transporter activity is integrated with the gene or the DNA according to any one of claims 8 to 10 in its chromosome, and stably shows high-affinity choline transporter activity (claim 46), and a cell having high-affinity choline transporter activity being obtainable by the preparing method of a cell having high-affinity choline transporter activity according to claim 45 or 46 (claim 47).

[0013]

The present invention also relates to a screening method of a promoter or a suppressor of high-affinity choline transporter activity characterized in measuring/evaluating high-affinity choline transporter activity of the protein having high-affinity choline transporter activity according to any one

of claims 14 to 22 in the presence of a subject material (claim 48), a screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in comprising the steps of: a cell membrane or a cell which expresses a protein having high-affinity choline transporter activity is cultivated in vitro in the presence of a subject material; the activity and/or the expression amount of a protein having high-affinity choline transporter activity in the cell membrane or the cell is measured/evaluated (claim 49), the screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to claim 49, wherein the cell membrane or the cell which expresses a protein having high-affinity choline transporter activity is the host cell containing an expression system which can express a protein having high-affinity choline transporter activity according to any one of claims 34 to 38, or is the cell having high-affinity choline transporter activity according to claim 47 (claim 50), the screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to any one of claims 48 to 50, wherein the protein having high-affinity choline transporter activity is a recombinant protein (claim 51), a screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in comprising the steps of: a cell obtained from the non-human animal according to any one of claims 39 to 44 is cultivated in vitro in the presence of a subject material; the activity and/or the expression amount of a protein having

high-affinity choline transporter activity in the cell is measured/evaluated (claim 52), a screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity (claim 53), a screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome, and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity (claim 54), a screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression characterized in administering a subject material to a non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome, and then evaluating the activity and/or the expression amount of a protein having high-affinity choline transporter activity in comparison with the case using wild-type non-human animal (claim 55), and the screening method of a promoter or a suppressor of high-affinity choline transporter activity, or of high-affinity choline transporter expression according to any one of claims 52 to 55, wherein the non-human animal is a mouse or a rat (claim 56).

[0014]

The present invention further relates to a material which promotes activity or expression of a protein having high-affinity choline transporter activity being obtainable by the screening method according to any one of claims 48 to 56 (claim 57), a material which suppresses activity or expression of a protein having high-affinity choline transporter activity being obtainable by the screening method according to any one of claims 48 to 56 (claim 58), a medical constituent characterized in being used for a medical treatment for a patient who needs elevation of the activity or enhancement of the expression of a high-affinity choline transporter, and containing the protein according to any one of claims 14 to 22, and/or the material which promotes activity or expression of a protein having high-affinity choline transporter activity according to claim 57 as an active component (claim 59), and a medical constituent characterized in being used for medical treatment for a patient who needs suppression of the activity or the expression of a high-affinity choline transporter, and containing the protein according to any one of claims 14 to 22, and/or the material which suppresses the activity or the expression of a protein having high-affinity choline transporter activity according to claim 58 as an active component (claim 60).

[0015]

The present invention still further relates to a diagnostic method for diseases relating to the expression or the activity of a high-affinity choline transporter characterized in comparing a DNA sequence encoding a high-affinity choline transporter in a sample to a DNA sequence encoding the protein according to claim 19 or 20 (claim 61), a diagnostic probe for Alzheimer's disease comprising a whole or a part of an antisense

strand of DNA or RNA encoding the protein according to claim 19 or 20 (claim 62), and a diagnostic drug for Alzheimer's disease characterized in containing the diagnostic probe according to claim 62 and/or the antibody according to any one of claims 28 to 33 (claim 63). [0015]

[0016]

[Mode for Carrying out the Invention]

The cDNA of nematode high-affinity choline transporter of the present invention, being described in Seq. ID No. 1, can be obtained by injecting each cRNA prepared from candidate full-length cDNAs, which are expected as a member of Na^+ -dependent transporter family according to *C. elegans* genome project, into oocytes of *Xenopus*, and examining the uptake of choline. The high-affinity uptake of choline in brain synaptosomes of mammals was completely inhibited by 1 μM hemicholinium-3 (HC3) ($K_i=10-100$ nM), while the low-affinity uptake of choline, which is distributed in every cells, was inhibited only by HC3 with higher concentration ($K_i=50$ μM). Therefore, the sensitivity to 1 μM HC3 can be used as criteria of high-affinity choline uptake during the process. For example, it is possible to confirm the identification, the expression, and the localization of an object gene from the candidate cDNA of a nematode (*C. elegans*) as follows.

[0017]

It has been found that cDNA corresponding to the gene expected as C48D1.3 promotes significant choline uptake, being inhibited by 1 μM HC3, in the high-affinity choline uptake process. Fig. 1 shows the result of [^3H] choline uptake of oocytes from *Xenopus* being injected with C48D1.3 cRNA or water. In Fig. 1, the closed and the open columns indicate choline uptake in the absence or the presence of 1 μM HC3 respectively, and each column

is shown by mean \pm SEM (n=6 to 8 oocytes). Fig. 2 shows the effect of Na⁺ on the choline uptake, and the closed columns indicate choline uptake measured in the standard solution ([Na⁺]=100 mM), the open columns indicate choline uptake in the absence of Na⁺ (Na⁺ was substituted with Li⁺). In addition, Fig. 3 shows the inhibition of choline uptake induced by HC3. Based on the above-mentioned Fig. 2 and 3, it is presumed that the uptake is Na⁺-dependent, and that K_i of HC3 is 50 nM. The cDNA clone was designated as cho-1 (high-affinity choline transporter-1).

[0018]

By comparing a base sequence of cDNA and that of genome, cho-1 gene was found to comprise 9 exons. A protein expected from a base sequence of cDNA of cho-1 includes 576 amino acid residues (see Fig. 4), and this protein, being represented by Seq. ID No. 2, can be constructed by a usual method. When the available data base was searched, the amino acid sequences of cho-1 showed weak, but significant homology to members of Na⁺-dependent glucose transporter family. Hydrophobic analysis and comparison to other transporters suggest that there is a twelve-transmembrane region (see Fig. 7).

[0019]

Then, in order to identify cells expressing cho-1 in the nervous system of a nematode (*C. elegans*), a gene of a green fluorescent protein (GFP) fused with a region 5.1kb upstream from cho-1 gene was introduced into a nematode, and distribution of neurons expressing cho-1::gfp was examined. A photograph of L1 larva possessing cho-1::gfp reporter DNA at the outside of chromosome is shown as Fig. 5 (scale bar; 50 μ m). In Fig. 5, the arrowhead indicates nerve ring. In the ventral nerve cord, GFP is expressed only in cholinergic motor nerve, however,

some of DA, DB nerve cells do not express GFP owing probably to deficiency of reporter DNA at the outside of chromosome. It supports the idea that cho-1 is a high-affinity choline transporter of the cholinergic neuron.

[0020]

The cDNA of rat high-affinity choline transporter of the present invention, being described in Seq. ID No. 3, can be prepared, for example, by a method comprising the steps of: paying attention to cho-1 homologous molecules of vertebrates and searching data base with amino acid sequences expected from cho-1, and identifying one candidate (GenBank accession number: AQ316435) in human genomic survey sequence (GSS); amplifying cDNA fragments from rat spinal cord cDNA by PCR with degenerate primers on the basis of homology of base sequences between the human genome DNA and cho-1; screening rat spinal cord cDNA library with this fragment, and a positive cDNA clone was obtained. A protein with 580 amino acid residues showing 51% identity and 70% similarity to cho-1 was expected from the base sequence of the longest reading frame (see Fig. 4). This rat cDNA clone was designated as CHT1. In Fig. 4, each amino acid sequence of rat CHT1 and nematode CHO-1 is shown, and the identical and the similar residues are indicated on a black ground and a gray ground respectively. The expected transmembrane region I-XII is underlined. This protein represented by Seq. ID No. 4 can be constructed by a usual method.

[0021]

The above-mentioned amino acid sequence of CHT1 is significantly homologous to members of Na⁺-dependent glucose transporter family (20 to 25%). The phylogenetic tree of Na⁺-dependent glucose transporter family made by

neighbor-joining method using a program CLUSTALW of National Institute of Genetics (Mishima, Japan) is shown in Fig. 6. In Fig. 6, the percentage of the identical amino acids, being contained in each protein, to rat CHT1 is shown on the right side. On the other hand, no homology was observed to a yeast choline transporter (J. Biol. Chem. 265, 15996-16003, 1990), a creatine transporter which had been originally reported as a high-affinity choline transporter (Biochem. Biophys. Res. Commun. 198, 637-645, 1994), and other neurotransmitter transporters.

[0022]

The expected topology of CHT1 is thought to be the same as that of nematode CHO-1 fundamentally. Fig. 7 shows the expected topology of rat CHT1. In Fig. 7, the closed circles indicate the identical residues, the shadowed circles indicate highly conserved residues, and open circles indicate nonsimilar residues. The offshoots indicate the expected glycosylation sites. P among the circles shows the expected parts of phosphorylation induced by protein kinase C.

[0023]

Next, the distribution of CHT1 mRNA expression was examined by Northern blot analysis and *in situ* hybridization. The expression of transcripts with the length of about 5 kb was confirmed by Northern blot analysis of various tissues of rats. Fig. 8 shows the result of Northern blot analysis of mRNA transcript of CHT1 in rat tissue, and the length of RNA standard (0.24 to 9.5 kb; GIBCO BRL) is exhibited on the left side. As shown in Fig. 8, an abundance of transcripts were confirmed in basal forebrain, brain stem and spinal cord, and a little of those were confirmed in corpus striatum. These tissues are known

to contain cholinergic neurons. On the other hand, no transcript was observed in other regions of the brain or in tissues of non-nervous systems.

[0024]

Consistent with these results, *in situ* hybridization confirmed the expression of CHT1 mRNA in cell groups of main cholinergic neurons including corpus striatum, cell population in basal forebrain and ventral horn in spinal cord. Fig. 9 and 10 (scale bar; 1 mm) show micrographs of sections in bright-field, which were hybridized with a cRNA probe of an antisense labeled by digoxigenin. These micrographs relate to *in situ* hybridization analysis of CHT1 transcripts in rat brain and spinal cord. Fig. 9 indicates that mRNA transcripts of CHT1 were detected in vertical and horizontal limbs of the diagonal band (VDB, HDB), medial septal nucleus (MS), caudate and putamen (Cpu), and olfactory tubercle (Tu). Fig. 10 indicates that the expression was observed in ventral horn (VH) in spinal cord. Further, the adjacent section hybridized with a probe of vesicle acetylcholine transporter showed essentially same distribution. This expression distribution is essentially same as the reported distribution of choline acetyl group transferase or vesicle acetylcholine transporter. These results show that the expression of CHT1 mRNA is limited to cholinergic neurons.

[0025]

Next, choline uptake of CHT1 was examined by using oocytes of *Xenopus*. The choline uptake of the oocytes injected with CHT1 cRNA was 2 times to 4 times more than that of controls injected with water. Fig. 11 shows the result of [³H] choline uptake of oocytes of *Xenopus* injected with CHT1 cRNA or water. In Fig. 11, the open and the closed columns respectively indicate choline

uptake in the standard solutions containing 100 mM NaCl or LiCl, and each column is shown by mean \pm SEM (n=6 to 8 oocytes). The effect of choline concentration on choline uptake is shown in Fig. 12. In Fig. 12, choline uptake of oocytes injected with water was subtracted from that of oocytes injected with cRNA in order to figure out CHT1-induced choline uptake, and the choline uptake was fitted to Michaelis-Menten curve. As shown in Fig. 12, choline uptake of CHT1 saturated when increasing choline concentration ($K_m=2.2 \pm 0.2 \mu\text{M}$, n=3).

[0026]

Then, the result of HC3-induced inhibition of choline uptake is shown in Fig. 13. Fig. 13 indicates that the K_m of endogenous choline uptake of control is higher than 10 μM and that choline uptake of CHT1 is completely inhibited by 0.1 μM HC3 ($K_i=2-3 \text{ nM}$), whereas 10 μM HC3 induced only slight inhibition in control. As shown in Fig. 14, ion-dependency of choline uptake of CHT1 was examined and found to be Cl^- -dependent as well as Na^+ -dependent. The closed and the open columns indicate choline uptake of oocytes injected with water and with cRNA respectively (100 mM NaCl in the standard solution is substituted with 100 mM of each salt) shown in the figure. These results indicate that CHT1 has the characteristics expected from high-affinity choline uptake in brain synaptosomes (high-affinity to choline, high sensitivity to HC3, and Na^+ - Cl^- -dependency) (J. Neurochem. 27, 93-99, 1976).

[0027]

In addition, [^3H] HC3 binding activity of membranes prepared from COS7 cells introduced with CHT1 cDNA and a vector (control) respectively was examined. The result is shown in Fig. 15. As Fig. 15 indicates, Na^+ -dependent [^3H] HC3 binding

was observed in a membrane of a cell where CHT1 was expressed, but not in a control membrane. Subsequently, a saturation analysis was conducted for specific [³H] HC3 binding. As shown in Fig. 16, equilibrium dissociation constant (K_d) was estimated to be 1.6±0.2 μM (n=3). This value was similar to that reported in brain synaptosomes (J. Neurochem. 60, 1191-1201, 1993, Life Sci. 35, 2335-2343, 1984, Brain Res. 348, 321-330, 1985). Further, displacement of specific [³H] HC3 binding by HC3, choline (Cho) and acetylcholine (Ach) was examined. Acetylcholine was measured in the presence of 1 μM physostigmine. The result is shown in Fig. 17. Fig. 17 indicates that specific [³H] HC3 binding was displaced when the concentration of choline was at least about 10 times lower than that of acetylcholine. These results show that CHT1 is a HC3 binding site as well as a high-affinity choline transporter.

[0028]

The cDNA of human high-affinity choline transporter of the present invention, being represented by Seq. ID No.5, can be prepared, for example, as follows: data base search was conducted with the amino acid sequence of nematode (*C. elegans*) CHO-1 to find a sequence of specific human genome DNA fragment having significant homology (R-107P12, a clone of human genomic survey sequence; GenBank accession number: AQ316435); a gene-specific primers for PCR were designed based on a base sequence of said DNA fragment; 5'-RACE (rapid amplification of cDNA ends) and 3'-RACE were conducted using Marathon-ReadyTM cDNA (Clontech) of human whole brain, together with an attached adapter primer; the obtained PCR product was cloned into a cloning vector for PCR, and a base sequence of inserted DNA was determined. In addition, an amino acid sequence expected from this DNA

sequence is represented by Seq. ID No. 6. A protein having human high-affinity choline transporter activity represented by said Seq. ID No. 6 can be constructed by a usual method on the basis of DNA sequence information shown in Seq. ID No. 5.

[0029]

The cDNA of mouse high-affinity choline transporter of the present invention, being represented by Seq. ID No.7, can be prepared, for example, as follows: data base search was conducted with the amino acid sequence of nematode (*C. elegans*) CHO-1 to find a sequence of specific human genome DNA fragment having significant homology (R-107P12, a clone of human genomic survey sequence; GenBank accession number: AQ316435); a gene-specific primer for PCR was designed based on a base sequence of said DNA fragment; 5'-RACE (rapid amplification of cDNA ends) and 3'-RACE were conducted using Marathon-ReadyTM cDNA (Clontech) of mouse whole brain, together with an attached adapter primer; the obtained PCR product was cloned into a cloning vector for PCR, and a base sequence of inserted DNA was determined. In addition, an amino acid sequence expected from this DNA sequence is represented by Seq. ID No. 8. A protein having mouse high-affinity choline transporter activity represented by said Seq. ID No. 8 can be constructed by a usual method on the basis of DNA sequence information shown in Seq. ID No. 7.

[0030]

Examples of a protein having high-affinity choline transporter activity of the present invention include a protein derived from natural materials and a recombinant protein. In addition to the ones represented by Seq. ID Nos. 2, 4, 6 and 8, which are specifically disclosed above, a protein comprising an amino acid sequence wherein one or a few amino acids are

deficient, substituted or added in amino acid sequences represented by Seq. ID Nos. 2, 4, 6 and 8, and having high-affinity choline transporter activity is also included. These proteins can be prepared by known methods. Further, examples of a gene or DNA encoding a protein having high-affinity choline transporter activity of the present invention include, in addition to the ones represented by Seq. ID Nos. 1, 3, 5 and 7, which are specifically disclosed above, a gene or DNA which encodes a protein comprising an amino acid sequence wherein one or a few amino acids are deficient, substituted or added in amino acid sequences represented by Seq. ID Nos. 2, 4, 6 and 8, and having high-affinity choline transporter activity, and DNA which encodes a protein hybridizing with said gene or DNA under a stringent condition and having high-affinity choline transporter activity. These genes and DNAs can be prepared by known methods.

[0031]

Cholinergic neurons play an extremely important role in learning and memory. The damage of these neurons correlates to severity of dementia. The rate-limiting step in acetylcholine synthesis is presumed to be the uptake of choline, and its activity is controlled by neural activity or various kinds of stimuli. In the brains of patients who suffer Alzheimer's disease, the hyperfunction of high-affinity choline uptake and of HC3 binding activity are observed (Trends Neurosci. 15, 117-122, 1992, Ann. NY Acad. Sci. 777, 197-204, 1996, J. Neurochem. 69, 2441-2451, 1997). Cloning of said gene or DNA encoding a protein having high-affinity choline transporter activity and said protein having high-affinity choline transporter activity is important for elucidating the molecular

mechanism of the high-affinity choline transporter and for developing new therapies for Alzheimer's disease.

[0032]

The fusion protein of the present invention means a substance constructed by binding a protein from a nematode, a rat, a human, a mouse, etc., which has high-affinity choline transporter activity, to a marker protein and/or a peptide tag. As the marker protein, any conventionally known marker protein can be used and the specific examples are alkaline phosphatase, Fc region of an antibody, HRP, and GFP. Conventionally known peptide tags, such as Myc tag, His tag, FLAG tag, GST tag, are exemplified as specific examples of the peptide tag of the present invention. Said fusion proteins can be constructed by a usual method, and are useful for the purification of a protein having high-affinity choline transporter activity utilizing the affinity between Ni-NTA and His tag, the detection of a protein having high-affinity choline transporter activity, the quantitation of an antibody to a protein having high-affinity choline transporter activity, and as a diagnostic marker for Alzheimer's disease, and an investigational reagent in the field concerned.

[0033]

As an antibody that specifically combines with a protein having high-affinity choline transporter activity of the present invention, an immunospecific antibody such as a monoclonal antibody, a polyclonal antibody, a chimeric antibody, a single stranded antibody, a humanized antibody and the like are concretely exemplified. Though these antibodies can be constructed by a usual method with the above-mentioned protein having high-affinity choline transporter activity as an antigen,

a monoclonal antibody is more preferable among them because of its specificity. Said antibody that specifically binds to a protein having high-affinity choline transporter activity, such as a monoclonal antibody or the like, is useful, for instance, for the diagnosis of Alzheimer's disease, and for elucidation of molecular mechanism of a high-affinity choline transporter.

[0034]

An antibody to a protein having high-affinity choline transporter activity is developed by administering fragments containing the protein having high-affinity choline transporter activity or its epitope, or cells that express said protein on the surface of the membrane, to animals (preferably excluding human) with usual protocol. For instance, a monoclonal antibody can be prepared by an arbitrary method that brings antibodies developed by cultured materials of continuous cell line, such as hybridoma method (Nature 256, 495-497, 1975), trioma method, human B-cell hybridoma method (Immunology Today 4, 72, 1983), and EBV-hybridoma method (MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

[0035]

In order to develop a single stranded antibody to the above-mentioned protein having high-affinity choline transporter activity of the present invention, the preparation method of single stranded antibodies (US Patent No. 4,946,778) can be applied. Further, in order to express a humanized antibody, it is possible to use transgenic mice, other mammalian animals or the like, and to isolate and identify the clones that express a protein having high-affinity choline transporter activity with the above-mentioned antibodies, and to purify the polypeptide by affinity chromatography. An antibody to a protein having

high-affinity choline transporter activity could be used, in particular, for the diagnosis and the medical treatment of Alzheimer's disease, and the like.

[0036]

This invention relates to a host cell which contains an expression system that can express said protein having high-affinity choline transporter activity. The gene that encodes a protein having high-affinity choline transporter activity can be introduced into a host cell by a number of methods described in many standard laboratory manuals such as by Davis et al. (BASIC METHODS IN MOLECULAR BIOLOGY, 1986), and by Sambrook et al. (MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y., 1989). Examples of those methods include calcium phosphate transfection, DEAE-dextran-mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection. Examples of the host cells include bacterial procaryotic cells such as Escherichia coli, Streptomyces, Bacillus subtilis, Streptococcus, Staphylococcus and the like; fungous cells such as yeast, Aspergillus and the like; insect cells such as drosophila S2, spodptera Sf9 and the like; and animal or plant cells such as L cells, CHO cells, COS cells, HeLa cells, C127 cells, BALB/c3T3 cells (including mutant strains deficient in dihydrofolate reductase, thymidine kinase or the like), BHK21 cells, HEK293 cells, Bowes melanoma cells and the like.

[0037]

As the expression system, any expression system that can express a protein having high-affinity choline transporter

activity in a host cell will suffice. Examples of the expression system include expression systems derived from chromosome, episome and virus, for example, vectors derived from bacterial plasmid, yeast plasmid, papovavirus like SV40, vaccinia virus, adenovirus, chicken pox virus, pseudorabies virus, or retrovirus, vectors derived from bacteriophage, transposon, and the combination of these, for instance, vectors derived from genetic factors of plasmid and of bacteriophage such as cosmid or phagemid. These expression systems may contain a regulatory sequence that acts not only as a promoter but also as a controller of expressions.

[0038]

A host cell that contains the above-mentioned expression system, cell membrane of said host cell, and a protein having high-affinity choline transporter activity which is obtainable by the cultivation of said host cell can be used in the screening method of the present invention as hereinafter described. For example, the method of F. Pietri-Rouxel et al. (Eur. J. Biochem., 247, 1174-1179, 1997) or the like can be used as the method to obtain cell membranes, and publicly known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxyapatite chromatography, and lectin chromatography, preferably high-speed liquid chromatography can be used to pick up said protein having high-affinity choline transporter activity from cell cultured material and purify it. As columns used for affinity chromatography, in particular, there are columns to which a protein antibody having anti-high-affinity choline transporter activity is bound, or in case that a normal peptide tag is added

to said high-affinity choline transporter, there are columns to which materials having affinity to the peptide tag are bound. Proteins having high-affinity choline transporter activity can be obtained by using these columns.

[0039]

In the present invention, said non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient on its chromosome means a non-human animal wherein a part or a whole of a gene encoding a protein having high-affinity choline transporter activity on chromosome is inactivated by gene mutation such as disruption, deficiency, substitution, etc. and function of expressing a protein having high-affinity choline transporter activity is lost. In addition, a non-human animal that overexpresses function of a gene that encodes a protein having high-affinity choline transporter activity on its chromosome means a non-human animal that produces larger amount of a protein having high-affinity choline transporter activity than a wild-type non-human animal does. Though specific examples of a non-human animal of the present invention include rodents, such as mice, rats and the like, a non-human animal of the present invention is not limited to these animals.

[0040]

Homozygous non-human animals generated according to Mendelian ratio include a deficient type or an overexpression type for a protein having high-affinity choline transporter activity, and their littermate wild-type, and it is possible to carry out precise comparative experiments in individual level by using the deficient types, the overexpression types and the littermate wild-types of these homozygous non-human animals at

the same time. Therefore, it is preferable to use animals of the same species, more preferably the littermates, as the wild-type non-human animals, in other words, the non-human animals being deficient in or overexpressing the function of a gene that encodes a protein having high-affinity choline transporter activity on their chromosome together in, for example, the screening hereinafter described in the present invention. The generating method of the non-human animals being deficient or overexpressing the function of a gene that encodes a protein having high-affinity choline transporter activity on their chromosome will be explained below, with an example of knockout mice and transgenic mice of a protein having high-affinity choline transporter activity.

[0041]

For example, a mouse being deficient in the function of a gene that encodes a protein having high-affinity choline transporter activity on its chromosome, in other words, a knockout mouse of a protein having high-affinity choline transporter activity on its chromosome can be constructed as follows. A gene that encodes a protein having high-affinity choline transporter activity is screened by using a gene fragment obtained from mouse gene library by a method like PCR. The screened gene that encodes a protein having high-affinity choline transporter activity is subcloned with a viral vector or the like, and specified by DNA sequencing. A target vector is constructed by substituting a whole or a part of a gene of this clone that encodes a protein having high-affinity choline transporter activity with pMC1 neo gene cassette or the like, and by introducing a diphtheria toxin A fragment (DT-A) gene, a herpes simplex virus thymidine kinase (HSV-tk) gene or other

such genes into 3'-terminal side.

[0042]

This constructed target vector is linearized and introduced into ES cells by electroporation or the like to induce homologous recombination. The ES cells wherein homologous recombination is induced by an antibiotic such as G418, ganciclovir (GANC) or the like are selected from the homologous recombinants. It is preferable to confirm whether the selected ES cells are the recombinants of the object by Southern blot or the like. A chimeric mouse is constructed by microinjecting a clone of the confirmed ES cells into a blastocyst of a mouse and then transplanting the blastocyst into a recipient mouse. A heterozygous mouse can be obtained by intercrossing the chimeric mouse with a wild-type mouse, and a knockout mouse of a protein having high-affinity choline transporter activity of the present invention can be constructed by intercrossing the heterozygous mice. It is possible to confirm whether a knockout mouse of a protein having high-affinity choline transporter activity is constructed, for example, by isolating RNA from the mouse obtained by said method and examining it by Northern blot analysis or the like, or by examining the expression of the mouse by Western blot analysis or the like.

[0043]

The transgenic mice of a protein having high-affinity choline transporter activity can be generated in following procedures. A transgene is constructed by fusing chicken β -actin, mouse neurofilament, SV40 or other such promoters, and rabbit β -globin, SV40 or other such poly A or introns with cDNA that encodes a protein having high-affinity choline transporter activity. The transgene is microinjected into the pronucleus

of a fertilized egg of a mouse, and the obtained egg cell is cultured, then transplanted to the oviduct of a recipient mouse. After rearing up the recipient animal, baby mice that have the above-mentioned cDNA are selected from the mice born from the recipient animal. Thus transgenic mice can be generated. The baby mouse that has cDNA can be selected by extracting crude DNA from a tail or the like of a mouse, then carrying out methods like dot hybridization using the introduced gene that encodes a protein having high-affinity choline transporter activity as a probe, PCR method using a specific primer, and the like.

[0044]

In addition, cells being useful for gene therapy of Alzheimer's disease and the like can be prepared by using a whole or a part of a gene or DNA that encodes a protein having high-affinity choline transporter activity of the present invention. As an example of a method for preparing these cells of the present invention, a method wherein a whole or a part of said gene or DNA of the present invention is introduced into a cell being deficient in the function of a gene that encodes a protein having high-affinity choline transporter activity on its chromosome by transfection or the like to obtain a cell having high-affinity choline transporter activity is exemplified. As the cell having high-affinity choline transporter activity, in particular, it is preferable to use a cell wherein said gene or DNA is integrated into a chromosome and high-affinity choline transporter activity is exhibited stably.

[0045]

By using the above-mentioned gene or DNA that encodes a protein having high-affinity choline transporter activity, a protein having high-affinity choline transporter activity, a

fusion protein created by combining a protein having high-affinity choline transporter activity and a marker protein and/or a peptide tag, an antibody to a protein having high-affinity choline transporter activity, a host cell which contains an expression system that can express a protein having high-affinity choline transporter activity, a cell having high-affinity choline transporter activity, or the like, it becomes possible to screen a pharmaceutical material useful for the treatment of symptoms as in Alzheimer's disease or the like, in other words, a material that promotes or suppresses the activity or the expression of a high-affinity choline transporter.

[0046]

Examples of the screening method of the present invention are: a method wherein the high-affinity choline transporter activity of the above-mentioned protein having high-affinity choline transporter activity of the present invention is measured/evaluated in the presence of a subject material; a method wherein a cell membrane or a cell which expresses a protein having high-affinity choline transporter activity of the present invention is cultivated in vitro in the presence of a subject material, and the activity and/or the expression amount of a protein having high-affinity choline transporter activity in the cell is measured/evaluated; and a method wherein a subject material is administered to said non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome and/or a wild-type non-human animal and then the activity and/or the expression amount of a protein having high-affinity choline transporter activity of the present

invention is measured/evaluated. As said cell membrane or said cell, a cell such as a primary cultured cell obtained from said non-human animal whose function of a gene encoding a protein having high-affinity choline transporter activity is deficient or overexpresses on its chromosome or a wild-type non-human animal etc., a host cell containing an expression system which can express a protein having high-affinity choline transporter activity of the present invention, a cell having high-affinity choline transporter activity of the present invention, and cell membranes of these cells can be specifically exemplified.

[0047]

The screening methods with said subject material and said protein having high-affinity choline transporter activity are now specifically explained together with examples, but the screening methods of the present invention are not limited to these examples. Cells expressing a protein having high-affinity choline transporter activity are cultured in the presence of a subject material, and the increase or the decrease of a protein having high-affinity choline transporter activity expressed on the cell surface after a certain period of cultivation can be immunochemically detected by ELISA or other such method with an antibody that specifically combines to a protein having high-affinity choline transporter activity of the present invention, or can be evaluated by using suppression or promotion of mRNA expression as an index. The mRNA can be detected by methods such as DNA chip, Northern hybridization or the like. Moreover, with a cell to which a gene wherein luciferase or other such reporter genes is linked to downstream of promoter of a gene that encodes high-affinity choline transporter is introduced, the suppression or the promotion of

the expression of a gene that encodes a protein having high-affinity choline transporter activity induced by a subject material can be detected by using the activity of said reporter gene as an index.

[0048]

The present invention further relates a medical constituent being used for medical treatment for a patient who needs promotion of the activity or the expression of a protein having high-affinity choline transporter, or a medical constituent being used for medical treatment for a patient who needs suppression of the activity or the expression of a protein having high-affinity choline transporter, wherein the material contains a protein having high-affinity choline transporter activity, a material which promotes the activity or the expression of a protein having high-affinity choline transporter activity, or a material which suppresses activity or expression of a protein having high-affinity choline transporter activity as an active component. As a protein having high-affinity choline transporter activity is involved in many biological functions including many pathological ones, it is expected that a compound that can stimulate a protein having high-affinity choline transporter activity and a compound being able to inhibit the function of said protein can be used as pharmaceuticals.

[0049]

As the material which promotes or suppresses the activity or the expression of a protein having high-affinity choline transporter activity, any material can be used as long as it binds to a protein having high-affinity choline transporter activity, or works on a signal transmitting molecule on upstream, and then promotes the activity or the expression of a protein

having high-affinity choline transporter activity or inhibits/antagonizes the activity or the expression of the protein by itself. Specific examples include an antibody, a ligand of a protein having high-affinity choline transporter activity, a fragment of said protein, and an oligonucleotide encoding said fragment, and these materials can be used as pharmaceuticals for treatment, prevention or the like of symptoms observed in the case of Alzheimer's disease or other such diseases, but use of them is not limited to the above examples.

[0050]

The present invention also relates to a diagnostic method for diseases relating to the activity or the expression of a protein having high-affinity choline transporter activity comprising a comparison of a DNA sequence encoding a protein having high-affinity choline transporter activity in a sample with a DNA sequence encoding a protein having high-affinity choline transporter activity of the present invention. The mutant type of DNA which encodes a protein having high-affinity choline transporter activity can be detected by finding gene-mutated individuals in DNA level, and this is useful for diagnosis of diseases caused by underexpression, overexpression or mutant expression of a protein having high-affinity choline transporter activity. Specific examples of a sample of said detection include cells of trial subjects, for example, genomic DNA, RNA or cDNA obtained from biopsy of blood, urine, saliva, tissue or the like, however said sample is not limited to these examples. It is also possible to use said sample being amplified by PCR or other such methods. Deficiency and insertion mutation of base sequences can be detected by the size change of the amplified product observed in comparison with normal genotype,

and point mutation can be identified by hybridizing amplified DNA with a labeled gene that encodes a protein having high-affinity choline transporter activity. Thus, diagnosis or judgement of symptoms observed in the case of Alzheimer's disease or other such diseases can be made by detecting the mutation of a gene that encodes a protein having high-affinity choline transporter activity.

[0051]

The present invention further relates to a diagnostic probe for diseases showing symptoms similar to those of Alzheimer's disease or the like comprising a whole or a part of an antisense chain of DNA or RNA encoding a protein having high-affinity choline transporter activity, and a diagnostic drug for diseases showing symptoms similar to those of Alzheimer's disease containing the diagnostic probe and/or an antibody which specifically binds to a protein having high-affinity choline transporter activity of the present invention. Said diagnostic probe is not limited in particular, as long as it comprises a whole or a part of an antisense chain of DNA (cDNA) or RNA (cRNA) encoding a protein having high-affinity choline transporter activity and being long enough to be a probe (at least 20 bases). In order to make a diagnostic drug for symptoms similar to those of Alzheimer's disease containing said probe and/or an antibody which specifically binds to a protein having high-affinity choline transporter activity of the present invention as active components, it is preferable to dissolve said probe into an appropriate buffer or sterilizing water for preventing said probe from decomposition. Further, it is also possible to diagnose diseases showing symptoms similar to those of Alzheimer's disease by methods using these diagnostic drugs, such as immunostaining

(Dev. Biol. 170, 207-222, 1995, J. Neurobiol. 29, 1-17, 1996), in situ hybridization (J. Neurobiol. 29, 1-17, 1996), in situ PCR or the like.

[0052]

Experimental methods or the like of the above-mentioned various experiments will now be explained in more detail below. (Cloning of high-affinity choline transporter cDNA)

The candidate cDNA of nematode high-affinity choline transporter was isolated from poly (A)+RNA of nematode mixture from various stages in the development by reverse transcription PCR and 3' RACE. MarathonTM cDNA Amplification Kit (Clontech) was used according to its protocol. A primer for sense direction of PCR was designed at a provisional translation initiating point of a predicted gene based on a DNA base sequence obtained from C. elegans genomic project. The amplified PCR product was subcloned into Nco I (smoothing) site and Not I site of a modified pSPUTK vector (Stratagene), and the base sequence of inserted DNA was determined. CHT1 cDNA of rat was isolated from rat spinal cord cDNA library by using GeneTrapper cDNA Positive Selection System (GIBCO Bio-Rad Laboratory: GIBCO BRL) according to its protocol. The primer used was designed from the base sequence of a cDNA fragment obtained by degenerated PCR. The obtained cDNA clones were analyzed. Among them, positive clones were selected and subcloned into pSPUTK vector and pcDNA3.1+ vector (Invitrogen Corporation).

[0053]

(Expression in oocytes of Xenopus)

In the presence of cap analog, cRNA was synthesized in vitro with SP6 or T7 RNA polymerase. 20 to 30 ng capped RNA was microinjected into oocytes (stage V to VI) of Xenopus. The

uptake was measured in basically same manner as described previously (Nature 360, 467-471, 1992). Two or three days after the injection of RNA, choline uptake was conducted for 30 to 60 min. with oocytes (6 to 8) in 0.75 ml standard solution (0.01 to 1 μ M [3 H]-choline, 100 mM NaCl, 2mM KCl, 1mM MgCl₂, 1mM CaCl₂, 10mM HEPES, 5mM Tris: pH 7.4). The oocytes completing uptake were solubilized with 10% SDS, and the amount of [3 H] was measured by a liquid scintillation counter.

[0054]

(GFC expression construct)

The transcriptional fusion construct of cho-1::gfp was constructed by PCR in same manner as described previously (Gene 212, 127-135, 1998). A gene that encodes a green fluorescent protein (GFP) located on downstream of a nuclear localization signal sequence (NLS) was inserted into a position 3 residues downstream of cho-1 translation initiating point so that the reading frame was fitted. NLS and gfp gene were amplified from pPD104.53 vector. In order to prepare 5.1 kb upstream region of cho-1 translation initiating point, a PCR primer being designed to encompass the first 3 amino acid residues of cho-1 was used. By the same method as previously described (EMBO J. 10, 3959-3970, 1991), rol-6 (sul006) marker and generated DNA were injected into gonads of a nematode simultaneously.

[0055]

(Northern blot analysis)

6 μ g poly(A)+RNA prepared from various tissues of rats was separated by formaldehyde-agarose electrophoresis, and transferred to a nylon membrane, then hybridized with CHT1 cDNA fragment being labeled with [32 P] by random prime method in hybridization solution (solution containing the final

concentration of 50% formamido, 5 × SSPE, 5 × Denhardt's solution, 0.5% SDS, 100 µg/ml salmon sperm DNA) at 42° C for 16 hours. The nylon membrane was washed under final condition (0.1 × SSPE, 0.1% SDS: 65° C), and then autoradiography was conducted for 7 days together with an enhancing screen.

[0056]

(In situ hybridization)

The transcript of an antisense labeled with digoxigenin was synthesized in vitro. Alkaline hydrolysis was repeated for the transcripts until their mean length was prepared to be 200 to 400 b. Cryostat sections of fresh frozen tissue (10 to 20 µm) were used. Hybridization was conducted with labeled cRNA probe (about 1 µg/ml) dissolved in 1 × Denhardt's solution [solution containing the final concentration of 50 mM Tris-HCl (pH 8.0), 2.5 mM EDTA, 0.3 M NaCl, 50% formamido, 10% dextran sulphate, 1 mg/ml E. coli tRNA] at 45° C for 20 hours. Then the sections were washed twice in 2 × SSC/50% formamido and once in 1 × SSC/50% formamido, at 45° C respectively. The hybridized probe was visualized by using anti-digoxigenin Fab fragment (Boehringer-Mannheim) and NBT/BCIP substrate. The sections were brought into reaction in substrate solution for 24 to 48 hours.

[0057]

(Binding assay)

[³H] hemicholinium-3 (HC3; 128Ci/mmol) was obtained from NEN Life Science Products. Either pcDNA3.1-CHT1 or pcDNA3.1 was transiently expressed in COS7 cells respectively. TransFast Reagent (Promega) was introduced and used according to the protocol. Membranes were prepared by following steps: homogenizing cells in 0.32 M sucrose; centrifuging the cells

for 1 hour at 200,000g; and suspending the precipitate. Binding assay was conducted in basically same manner as described previously. Specific binding amount was calculated by subtracting non-specific binding amount determined in the presence of 10 μ M HC3 from the whole binding amount. The Kd value was figured out by analyzing specific [3 H] HC3 binding amount from data of saturation binding assay with nonlinear approximation.

[0058]

[Effect of the Invention]

The present invention makes it possible to provide a protein having high-affinity choline transporter activity, which is physiologically important, and gene DNA encoding said protein. In addition, by using the said protein and gene DNA, it becomes possible to screen materials being useful for prevention or treatment of Alzheimer's disease, and to prepare cells being useful for gene therapy.

[0059]

[Sequence Listing]

SEQUENCE LISTING

<110> JAPAN SCIENCE AND TECHNOLOGY CORPORATION

<120> High-Affinity Choline Transporter

<130> A011P15

<140>

<141>

<150> JP P1999-240642

<151> 1999-08-27

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 1731

<212> DNA

<213> Caenorhabditis elegans

<220>

<221> CDS

<222> (1)..(1731)

<400> 1

atg gcc gac tta ttg ggt atc gtg gcc att gtg ttc ttc tac gtg ctc 48

Met Ala Asp Leu Leu Gly Ile Val Ala Ile Val Phe Phe Tyr Val Leu

1 5 10 15

att ctt gtc gtt gga ata tgg gcg ggt aga aaa tcg aaa agt tca aaa 96

Ile Leu Val Val Gly Ile Trp Ala Gly Arg Lys Ser Lys Ser Ser Lys

20 25 30

gag ctt gaa tca gaa gcc ggc gcg gcg acg gaa gag gtg atg tta gct 144

Glu Leu Glu Ser Glu Ala Gly Ala Ala Thr Glu Glu Val Met Leu Ala

35 40 45

ggg aga aac atc gga act ctt gtc gga att ttc aca atg act gcc acg 192

aca ttc acc ggt gga tac tat gca gtc gcg tac act gac gtc gtt caa 576
 Thr Phe Thr Gly Gly Tyr Tyr Ala Val Ala Tyr Thr Asp Val Val Gln
 180 185 190

cia ttt tgc att ttc gtc ggt ttg tgg gtt tgc gtg ccg gcg gct atg 624
 Leu Phe Cys Ile Phe Val Gly Leu Trp Val Cys Val Pro Ala Ala Met
 195 200 205

gtg cat gat ggt gcg aag gat att tcc agg aat gca ggc gac tgg att 672
 Val His Asp Gly Ala Lys Asp Ile Ser Arg Asn Ala Gly Asp Trp Ile
 210 215 220

gga gag att gga gga ttc aaa gaa aca tct ctc tgg att gat tgc atg 720
 Gly Glu Ile Gly Gly Phe Lys Glu Thr Ser Leu Trp Ile Asp Cys Met
 225 230 235 240

ctt ctc ctt gtc ttt gga gga att cca tgg caa gtg tac ttc caa aga 768
 Leu Leu Leu Val Phe Gly Gly Ile Pro Trp Gln Val Tyr Phe Gln Arg
 245 250 255

gtt ctc tcc tca aaa act gct cat gga gca cag acg ttg tgc ttt gtg 816
 Val Leu Ser Ser Lys Thr Ala His Gly Ala Gln Thr Leu Ser Phe Val
 260 265 270

gcg ggc gtc gga tgc att ctc atg gcg att cca cca gcg ttg atc ggt 864
 Ala Gly Val Gly Cys Ile Leu Met Ala Ile Pro Pro Ala Leu Ile Gly
 275 280 285

gca att gcc agg aac aca gac tgg aga atg act gat tat tcc cca tgg 912

Ala Ile Ala Arg Asn Thr Asp Trp Arg Met Thr Asp Tyr Ser Pro Trp

290

295

300

aac aat gga act aag gtc gaa tcg att cca ccg gat aag aga aac atg 960

Asn Asn Gly Thr Lys Val Glu Ser Ile Pro Pro Asp Lys Arg Asn Met

305

310

315

320

gtg gtc ccg ttg gta ttc cag tat ctt acg cca aga tgg gtc gcc ttt 1008

Val Val Pro Leu Val Phe Gln Tyr Leu Thr Pro Arg Trp Val Ala Phe

325

330

335

att gga ctc ggc gca gtg tcg gct gct gta atg tca tct gca gat tca 1056

Ile Gly Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser

340

345

350

tct gta cta tca gca gca tca atg ttt gct cac aac atc tgg aag ctc 1104

Ser Val Leu Ser Ala Ala Ser Met Phe Ala His Asn Ile Trp Lys Leu

355

360

365

aca att cgc cct cac gcg tct gaa aaa gaa gtg ata att gtg atg aga 1152

Thr Ile Arg Pro His Ala Ser Glu Lys Glu Val Ile Ile Val Met Arg

370

375

380

ata gcc atc atc tgt gtt ggt atc atg gca acc atc atg gca ctt acc 1200

Ile Ala Ile Ile Cys Val Gly Ile Met Ala Thr Ile Met Ala Leu Thr

385

390

395

400

att caa tcc atc tat ggg ctt tgg tat ctt tgt gca gat ttg gtc tac 1248

Ile Gln Ser Ile Tyr Gly Leu Trp Tyr Leu Cys Ala Asp Leu Val Tyr

405

410

415

gic ata ctc ttc cct caa cia tta tgt gtt gta tat atg cca cgt agc 1296
 Val Ile Leu Phe Pro Gln Leu Leu Cys Val Val Tyr Met Pro Arg Ser
 420 425 430

aat acg tat ggc tca ttg gct ggc tat gca gtc ggt ctt gtg ctc cgt 1344
 Asn Thr Tyr Gly Ser Leu Ala Gly Tyr Ala Val Gly Leu Val Leu Arg
 435 440 445

ttg att gga ggc gag cca ctt gta tgc ctg cca gcg ttc ttc cat tat 1392
 Leu Ile Gly Gly Glu Pro Leu Val Ser Leu Pro Ala Phe Phe His Tyr
 450 455 460

cca atg tat acg gat ggg gta cag tat ttc cca ttc agg aca act gct 1440
 Pro Met Tyr Thr Asp Gly Val Gln Tyr Phe Pro Phe Arg Thr Thr Ala
 465 470 475 480

atg tta tct tca atg gct act atc tac att gta tca ata caa tcg gag 1488
 Met Leu Ser Ser Met Ala Thr Ile Tyr Ile Val Ser Ile Gln Ser Glu
 485 490 495

aag ctg ttc aaa tcg gga cgt ttg tct ccg gag tgg gac gta atg ggt 1536
 Lys Leu Phe Lys Ser Gly Arg Leu Ser Pro Glu Trp Asp Val Met Gly
 500 505 510

tgt gta gtg aat att ccg ata gat cat gta ccc ctt ccg tca gat gta 1584
 Cys Val Val Asn Ile Pro Ile Asp His Val Pro Leu Pro Ser Asp Val
 515 520 525

tcg ttt gct gtt agt agt gag acc ttg aat atg aag gct cca aac gga 1632

Ser Phe Ala Val Ser Ser Glu Thr Leu Asn Met Lys Ala Pro Asn Gly

530

535

540

aca ccg gct cca gta cat ccg aac caa cag ccg tct gat gaa aat aca 1680

Thr Pro Ala Pro Val His Pro Asn Gln Gln Pro Ser Asp Glu Asn Thr

545

550

555

560

tta tta cat cca tat tcg gac caa agt tat tat tcc aca aat agc aat 1728

Leu Leu His Pro Tyr Ser Asp Gln Ser Tyr Tyr Ser Thr Asn Ser Asn

565

570

575

taa

1731

<210> 2

<211> 576

<212> PRT

<213> *Caenorhabditis elegans*

<400> 2

Met Ala Asp Leu Leu Gly Ile Val Ala Ile Val Phe Phe Tyr Val Leu

1

5

10

15

Ile Leu Val Val Gly Ile Trp Ala Gly Arg Lys Ser Lys Ser Ser Lys

20

25

30

Glu Leu Glu Ser Glu Ala Gly Ala Ala Thr Glu Glu Val Met Leu Ala

35

40

45

Gly Arg Asn Ile Gly Thr Leu Val Gly Ile Phe Thr Met Thr Ala Thr

50

55

60

Trp Val Gly Gly Ala Tyr Ile Asn Gly Thr Ala Glu Ala Leu Tyr Asn

65

70

75

80

Gly Gly Leu Leu Gly Cys Gln Ala Pro Val Gly Tyr Ala Ile Ser Leu

85

90

95

Val Met Gly Gly Leu Leu Phe Ala Lys Lys Met Arg Glu Glu Gly Tyr

100

105

110

Ile Thr Met Leu Asp Pro Phe Gln His Lys Tyr Gly Gln Arg Ile Gly

115

120

125

Gly Leu Met Tyr Val Pro Ala Leu Leu Gly Glu Thr Phe Trp Thr Ala

130

135

140

Ala Ile Leu Ser Ala Leu Gly Ala Thr Leu Ser Val Ile Leu Gly Ile

145

150

155

160

Asp Met Asn Ala Ser Val Thr Leu Ser Ala Cys Ile Ala Val Phe Tyr

165

170

175

Thr Phe Thr Gly Gly Tyr Tyr Ala Val Ala Tyr Thr Asp Val Val Gln

180

185

190

Leu Phe Cys Ile Phe Val Gly Leu Trp Val Cys Val Pro Ala Ala Met

195

200

205

Val His Asp Gly Ala Lys Asp Ile Ser Arg Asn Ala Gly Asp Trp Ile
210 215 220

Gly Glu Ile Gly Gly Phe Lys Glu Thr Ser Leu Trp Ile Asp Cys Met
225 230 235 240

Leu Leu Leu Val Phe Gly Gly Ile Pro Trp Gln Val Tyr Phe Gln Arg
245 250 255

Val Leu Ser Ser Lys Thr Ala His Gly Ala Gln Thr Leu Ser Phe Val
260 265 270

Ala Gly Val Gly Cys Ile Leu Met Ala Ile Pro Pro Ala Leu Ile Gly
275 280 285

Ala Ile Ala Arg Asn Thr Asp Trp Arg Met Thr Asp Tyr Ser Pro Trp
290 295 300

Asn Asn Gly Thr Lys Val Glu Ser Ile Pro Pro Asp Lys Arg Asn Met
305 310 315 320

Val Val Pro Leu Val Phe Gln Tyr Leu Thr Pro Arg Trp Val Ala Phe
325 330 335

Ile Gly Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser
340 345 350

Ser Val Leu Ser Ala Ala Ser Met Phe Ala His Asn Ile Trp Lys Leu
355 360 365

Thr Ile Arg Pro His Ala Ser Glu Lys Glu Val Ile Ile Val Met Arg

370

375

380

Ile Ala Ile Ile Cys Val Gly Ile Met Ala Thr Ile Met Ala Leu Thr

385

390

395

400

Ile Gln Ser Ile Tyr Gly Leu Trp Tyr Leu Cys Ala Asp Leu Val Tyr

405

410

415

Val Ile Leu Phe Pro Gln Leu Leu Cys Val Val Tyr Met Pro Arg Ser

420

425

430

Asn Thr Tyr Gly Ser Leu Ala Gly Tyr Ala Val Gly Leu Val Leu Arg

435

440

445

Leu Ile Gly Gly Glu Pro Leu Val Ser Leu Pro Ala Phe Phe His Tyr

450

455

460

Pro Met Tyr Thr Asp Gly Val Gln Tyr Phe Pro Phe Arg Thr Thr Ala

465

470

475

480

Met Leu Ser Ser Met Ala Thr Ile Tyr Ile Val Ser Ile Gln Ser Glu

485

490

495

Lys Leu Phe Lys Ser Gly Arg Leu Ser Pro Glu Trp Asp Val Met Gly

500

505

510

Cys Val Val Asn Ile Pro Ile Asp His Val Pro Leu Pro Ser Asp Val

515

520

525

Ser Phe Ala Val Ser Ser Glu Thr Leu Asn Met Lys Ala Pro Asn Gly

530

535

540

Thr Pro Ala Pro Val His Pro Asn Gln Gln Pro Ser Asp Glu Asn Thr

545

550

555

560

Leu Leu His Pro Tyr Ser Asp Gln Ser Tyr Tyr Ser Thr Asn Ser Asn

565

570

575

<210> 3

<211> 1743

<212> DNA

<213> Rat

<220>

<221> CDS

<222> (1)..(1743)

<400> 3

atg cct ttc cat gta gaa gga cta gta gcg att atc ctg ttc tac ctt 48

Met Pro Phe His Val Glu Gly Leu Val Ala Ile Ile Leu Phe Tyr Leu

1

5

10

15

ctt ata ttt ctg gtt gga ata tgg gct gca tgg aaa acc aaa aac agc 96

Leu Ile Phe Leu Val Gly Ile Trp Ala Ala Trp Lys Thr Lys Asn Ser

20

25

30

ggt aat gca gaa gaa cgc agc gaa gcc atc ata gtt ggg ggc cga gac 144

Gly Asn Ala Glu Glu Arg Ser Glu Ala Ile Ile Val Gly Gly Arg Asp

35	40	45	
att ggt ttg ttg gtt ggt ggt ttt acc atg aca gcc acc tgg gtt gga			192
Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly			
50	55	60	
gga ggt tac atc aac ggg aca gct gaa gca gtt tat ggg cca ggt tgt			240
Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Gly Pro Gly Cys			
65	70	75	80
ggt cta gct tgg gct cag gca ccc att gga tat tct ctg agt ctg att			288
Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile			
	85	90	95
tta ggt ggc ctg ttt ttt gca aaa cct atg cgt tcc aag gga tat gtg			336
Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val			
100	105	110	
act atg tta gac ccg ttt caa cag atc tat gga aag cgc atg ggt ggg			384
Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly			
115	120	125	
ctg ctg ttc atc cct gca ctg atg gga gag atg ttc tgg gct gca gca			432
Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala			
130	135	140	
att ttc tct gca tta ggg gct acc atc agc gta atc att gat gig gat			480
Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp			
145	150	155	160

gig aac ata tgc gtc att gtc tcc gca ctc att gcc att ctt tat acc 528

Val Asn Ile Ser Val Ile Val Ser Ala Leu Ile Ala Ile Leu Tyr Thr

165

170

175

ctc gtg gga ggg ctc tac tct gtg gca tat act gat gtt gta cag cta 576

Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu

180

185

190

ttc tgc att ttt ata gga ttg tgg atc agt gtc cca ttt gcc ctg tca 624

Phe Cys Ile Phe Ile Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser

195

200

205

cat cct gca gtc acc gac att gga ttc act gct gtg cat gct aaa tac 672

His Pro Ala Val Thr Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr

210

215

220

cag agt ccc tgg ctg gga acc att gaa tca gtt gaa gtc tac acc tgg 720

Gln Ser Pro Trp Leu Gly Thr Ile Glu Ser Val Glu Val Tyr Thr Trp

225

230

235

240

ctt gat aat ttt ctg ttg ttg atg ctg ggt gga ata cca tgg caa gcc 768

Leu Asp Asn Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala

245

250

255

tac ttc cag agg gtc ctc tct tca tgc tca gcg acc tat gct cag gtg 816

Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val

260

265

270

ctg tcc ttc ctg gca gct ttt ggg tgc ctg gtg atg gct cta cca gcc 864

Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Leu Pro Ala

275	280	285	
att tgc att ggg gcc att gga gcc tcc aca gac tgg aac caa act gca 912			
Ile Cys Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala			
290	295	300	
tat ggg ttt cca gat ccc aag acc aag gag gaa gca gac atg att ctc 960			
Tyr Gly Phe Pro Asp Pro Lys Thr Lys Glu Glu Ala Asp Met Ile Leu			
305	310	315	320
ccg att gtt cta cag tac ctc tgc cct gtg tac att tcc ttc ttt ggg 1008			
Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly			
	325	330	335
ctt ggt gct gtt tct gct gct gtc atg tcc tcg gct gac tca tcc atc 1056			
Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile			
	340	345	350
cta tca gca agt tcc atg ttt gct cgg aat atc tac cag ctt tcc ttc 1104			
Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe			
	355	360	365
aga caa aat gca tca gac aag gaa att gtg tgg gtc atg agg atc act 1152			
Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr			
370	375	380	
gtg ttt gtg ttt gga gca tct gca aca gcc atg gcc ttg ctc acg aag 1200			
Val Phe Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys			
385	390	395	400

act gtg tat ggg ctc tgg tac ctg agc tct gac ctt gtc tac atc atc 1248
 Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Ile

405 410 415

atc ttc cca cag ctg ctc tgt gta ctc ttc atc aaa gga acc aac act 1296
 Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Ile Lys Gly Thr Asn Thr

420 425 430

tat ggg gca gtt gct ggt tat att ttt gga ctt ttc ctg aga att acc 1344
 Tyr Gly Ala Val Ala Gly Tyr Ile Phe Gly Leu Phe Leu Arg Ile Thr

435 440 445

gga gga gag cca tat cta tac ttg cag ccc tta atc ttc tac cct ggt 1392
 Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly

450 455 460

tat tac cct gac aag aat ggt ata tac aat cag agg ttc cca ttt aaa 1440
 Tyr Tyr Pro Asp Lys Asn Gly Ile Tyr Asn Gln Arg Phe Pro Phe Lys

465 470 475 480

act ctc tcc atg gtt acc tca ttc ttt acc aac att tgt gtt tcc tat 1488
 Thr Leu Ser Met Val Thr Ser Phe Phe Thr Asn Ile Cys Val Ser Tyr

485 490 495

cta gcc aag tat cta ttt gaa agt gga acc ttg cct cca aaa tta gat 1536
 Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp

500 505 510

ata ttt gat gct gtt gtc tca agg cac agt gaa gag aac atg gac aag 1584
 Ile Phe Asp Ala Val Val Ser Arg His Ser Glu Glu Asn Met Asp Lys

515 520 525
acc att cta gtc aga aat gaa aac atc aaa tta aat gaa ctt gca cct 1632
Thr Ile Leu Val Arg Asn Glu Asn Ile Lys Leu Asn Glu Leu Ala Pro

530 535 540
gta aag cct cga cag agc cta acc ctc agt tca act ttc acc aat aaa 1680
Val Lys Pro Arg Gln Ser Leu Thr Leu Ser Ser Thr Phe Thr Asn Lys
545 550 555 560

gag gct ctc ctt gat gtt gat tcc agt cca gag gga tct ggg act gaa 1728
Glu Ala Leu Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu
565 570 575

gat aac tta caa tga 1743
Asp Asn Leu Gln
580

<210> 4
<211> 580
<212> PRT
<213> Rat

<400> 4
Met Pro Phe His Val Glu Gly Leu Val Ala Ile Ile Leu Phe Tyr Leu
1 5 10 15
Leu Ile Phe Leu Val Gly Ile Trp Ala Ala Trp Lys Thr Lys Asn Ser
20 25 30

Gly Asn Ala Glu Glu Arg Ser Glu Ala Ile Ile Val Gly Gly Arg Asp

35

40

45

Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly

50

55

60

Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Gly Pro Gly Cys

65

70

75

80

Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile

85

90

95

Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val

100

105

110

Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly

115

120

125

Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala

130

135

140

Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp

145

150

155

160

Val Asn Ile Ser Val Ile Val Ser Ala Leu Ile Ala Ile Leu Tyr Thr

165

170

175

Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu

180

185

190

Phe Cys Ile Phe Ile Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser
195 200 205

His Pro Ala Val Thr Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr
210 215 220

Gln Ser Pro Trp Leu Gly Thr Ile Glu Ser Val Glu Val Tyr Thr Trp
225 230 235 240

Leu Asp Asn Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala
245 250 255

Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val
260 265 270

Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Leu Pro Ala
275 280 285

Ile Cys Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala
290 295 300

Tyr Gly Phe Pro Asp Pro Lys Thr Lys Glu Glu Ala Asp Met Ile Leu
305 310 315 320

Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly
325 330 335

Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile
340 345 350

Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe
355 360 365

Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr
370 375 380

Val Phe Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys
385 390 395 400

Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Ile
405 410 415

Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Ile Lys Gly Thr Asn Thr
420 425 430

Tyr Gly Ala Val Ala Gly Tyr Ile Phe Gly Leu Phe Leu Arg Ile Thr
435 440 445

Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly
450 455 460

Tyr Tyr Pro Asp Lys Asn Gly Ile Tyr Asn Gln Arg Phe Pro Phe Lys
465 470 475 480

Thr Leu Ser Met Val Thr Ser Phe Phe Thr Asn Ile Cys Val Ser Tyr
485 490 495

Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp
500 505 510

Ile Phe Asp Ala Val Val Ser Arg His Ser Glu Glu Asn Met Asp Lys
515 520 525

Thr Ile Leu Val Arg Asn Glu Asn Ile Lys Leu Asn Glu Leu Ala Pro
530 535 540

Val Lys Pro Arg Gln Ser Leu Thr Leu Ser Ser Thr Phe Thr Asn Lys
545 550 555 560

Glu Ala Leu Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu
565 570 575

Asp Asn Leu Gln
580

<210> 5

<211> 1743

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1743)

<400> 5

atg gct ttc cat gtg gaa gga ctg ata gct atc atc gtg ttc tac ctt 48

Met Ala Phe His Val Glu Gly Leu Ile Ala Ile Ile Val Phe Tyr Leu

1

5

10

15

cta att ttg ctg gtt gga ata tgg gct gcc tgg aga acc aaa aac agt	96
Leu Ile Leu Leu Val Gly Ile Trp Ala Ala Trp Arg Thr Lys Asn Ser	
20 25 30	
ggc agc gca gaa gag cgc agc gaa gcc atc ala gtt ggt ggc cga gat	144
Gly Ser Ala Glu Glu Arg Ser Glu Ala Ile Ile Val Gly Gly Arg Asp	
35 40 45	
att ggt tta ttg gtt ggt gga ttt acc atg aca gct acc tgg gtc gga	192
Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly	
50 55 60	
gga ggg tat atc aat ggc aca gct gaa gca gtt tat gta cca ggt tat	240
Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Val Pro Gly Tyr	
65 70 75 80	
ggc cta gct tgg gct cag gca cca att gga tat tct ctt agt ctg att	288
Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile	
85 90 95	
tta ggt ggc ctg ttc ttt gca aaa cct atg cgt tca aag ggg tat gtg	336
Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val	
100 105 110	
acc atg tta gac ccg ttt cag caa atc tat gga aaa cgc atg ggc gga	384
Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly	
115 120 125	
ctc ctg ttt att cct gca ctg atg gga gaa atg ttc tgg gct gca gca	432

Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala
 130 135 140

att ttc tct gct ttg gga gcc acc atc agc gtg atc atc gat gtg gat 480
 Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp
 145 150 155 160

atg cac att tct gtc atc atc tct gca ctc att gcc act ctg tac aca 528
 Met His Ile Ser Val Ile Ile Ser Ala Leu Ile Ala Thr Leu Tyr Thr
 165 170 175

ctg gtg gga ggg ctc tat tct gtg gcc tac act gat gtc gtt cag ctc 576
 Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu
 180 185 190

ttt tgc att ttt gta ggg ctg tgg atc agc gtc ccc ttt gca ttg tca 624
 Phe Cys Ile Phe Val Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser
 195 200 205

cat cct gca gtc gca gac atc ggg ttc act gct gtg cat gcc aaa tac 672
 His Pro Ala Val Ala Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr
 210 215 220

caa aag ccg tgg ctg gga act gtt gac tca tct gaa gtc tac tct tgg 720
 Gln Lys Pro Trp Leu Gly Thr Val Asp Ser Ser Glu Val Tyr Ser Trp
 225 230 235 240

ctt gat agt ttt ctg ttg ttg atg ctg ggt gga atc cca tgg caa gca 768
 Leu Asp Ser Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala
 245 250 255

tac ttt cag agg gtt ctc tct tct tcc tca gcc acc tat gct caa gtg 816
Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val
260 265 270

ctg tcc ttc ctg gca gct ttc ggg tgc ctg gtg atg gcc atc cca gcc 864
Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Ile Pro Ala
275 280 285

ata ctc att ggg gcc att gga gca tca aca gac tgg aac cag act gca 912
Ile Leu Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala
290 295 300

tat ggg ctt cca gat ccc aag act aca gaa gag gca gac atg att tta 960
Tyr Gly Leu Pro Asp Pro Lys Thr Thr Glu Glu Ala Asp Met Ile Leu
305 310 315 320

cca att gtt ctg cag tat ctc tgc cct gtg tat att tct ttc ttt ggt 1008
Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly
325 330 335

ctt ggt gca gtt tct gct gct gtt atg tca tca gca gat tct tcc atc 1056
Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile
340 345 350

tig tca gca agt tcc atg ttt gca cgg aac atc tac cag ctt tcc ttc 1104
Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe
355 360 365

aga caa aat gct tgc gac aaa gaa atc gtt tgg gtt atg cga atc aca 1152

Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr	
370 375 380	
gtg ttt gtg ttt gga gca tct gca aca gcc atg gcc ttg ctg acg aaa	1200
Val Phe Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys	
385 390 395 400	
act gtg tat ggg ctc tgg tac ctc agt tct gac ctt gtt tac atc gtt	1248
Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Val	
405 410 415	
atc ttc ccc cag ctg ctt tgt gta ctc ttt gtt aag gga acc aac acc	1296
Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Val Lys Gly Thr Asn Thr	
420 425 430	
tat ggg gcc gtg gca ggt tat gtt tct ggc ctc ttc ctg aga ata act	1344
Tyr Gly Ala Val Ala Gly Tyr Val Ser Gly Leu Phe Leu Arg Ile Thr	
435 440 445	
gga ggg gag cca tat ctg tat ctt cag ccc ttg atc ttc tac cct ggc	1392
Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly	
450 455 460	
tat tac cct gat gat aat ggt ata tat aat cag aaa ttt cca ttt aaa	1440
Tyr Tyr Pro Asp Asp Asn Gly Ile Tyr Asn Gln Lys Phe Pro Phe Lys	
465 470 475 480	
aca ctt gcc atg gtt aca tca ttc tta acc aac att tgc atc tcc tat	1488
Thr Leu Ala Met Val Thr Ser Phe Leu Thr Asn Ile Cys Ile Ser Tyr	
485 490 495	

cta gcc aag tat cta ttt gaa agt gga acc ttg cca cct aaa tta gat 1536
 Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp

500 505 510

gta ttt gat gct gtt gtt gca aga cac agt gaa gaa aac atg gat aag 1584
 Val Phe Asp Ala Val Val Ala Arg His Ser Glu Glu Asn Met Asp Lys

515 520 525

aca att ctt gtc aaa aat gaa aat att aaa tta gat gaa ctt gca ctt 1632
 Thr Ile Leu Val Lys Asn Glu Asn Ile Lys Leu Asp Glu Leu Ala Leu

530 535 540

gtg aag cca cga cag agc atg acc ctc agc tca act ttc acc aat aaa 1680
 Val Lys Pro Arg Gln Ser Met Thr Leu Ser Ser Thr Phe Thr Asn Lys

545 550 555 560

gag gcc ttc ctt gat gtt gat tcc agt cca gaa ggg tct ggg act gaa 1728
 Glu Ala Phe Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu

565 570 575

gat aat tta cag tga 1743

Asp Asn Leu Gln

580

<210> 6

<211> 580

<212> PRT

<213> Homo sapiens

<400> 6

Met Ala Phe His Val Glu Gly Leu Ile Ala Ile Ile Val Phe Tyr Leu

1 5 10 15

Leu Ile Leu Leu Val Gly Ile Trp Ala Ala Trp Arg Thr Lys Asn Ser

20 25 30

Gly Ser Ala Glu Glu Arg Ser Glu Ala Ile Ile Val Gly Gly Arg Asp

35 40 45

Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly

50 55 60

Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Val Pro Gly Tyr

65 70 75 80

Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile

85 90 95

Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val

100 105 110

Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly

115 120 125

Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala

130 135 140

Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp

145 150 155 160

Met His Ile Ser Val Ile Ile Ser Ala Leu Ile Ala Thr Leu Tyr Thr

165 170 175

Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu

180 185 190

Phe Cys Ile Phe Val Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser

195 200 205

His Pro Ala Val Ala Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr

210 215 220

Gln Lys Pro Trp Leu Gly Thr Val Asp Ser Ser Glu Val Tyr Ser Trp

225 230 235 240

Leu Asp Ser Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala

245 250 255

Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val

260 265 270

Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Ile Pro Ala

275 280 285

Ile Leu Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala

290 295 300

Tyr Gly Leu Pro Asp Pro Lys Thr Thr Glu Glu Ala Asp Met Ile Leu

305 310 315 320

Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly

325 330 335

Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile

340 345 350

Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe

355 360 365

Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr

370 375 380

Val Phe Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys

385 390 395 400

Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Val

405 410 415

Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Val Lys Gly Thr Asn Thr

420 425 430

Tyr Gly Ala Val Ala Gly Tyr Val Ser Gly Leu Phe Leu Arg Ile Thr

435 440 445

Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly

450 455 460

Tyr Tyr Pro Asp Asp Asn Gly Ile Tyr Asn Gln Lys Phe Pro Phe Lys

465 470 475 480

Thr Leu Ala Met Val Thr Ser Phe Leu Thr Asn Ile Cys Ile Ser Tyr

485 490 495

Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp

500 505 510

Val Phe Asp Ala Val Val Ala Arg His Ser Glu Glu Asn Met Asp Lys

515 520 525

Thr Ile Leu Val Lys Asn Glu Asn Ile Lys Leu Asp Glu Leu Ala Leu

530 535 540

Val Lys Pro Arg Gln Ser Met Thr Leu Ser Ser Thr Phe Thr Asn Lys

545 550 555 560

Glu Ala Phe Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu

565 570 575

Asp Asn Leu Gln

580

<210> 7

<211> 1743

<212> DNA

<213> Mouse

<220>

<221> CDS

<222> (1)..(1743)

<400> 7

atg tct ttc cac gta gaa gga ctg gta gct att atc ctc ttc tac ctc 48

Met Ser Phe His Val Glu Gly Leu Val Ala Ile Ile Leu Phe Tyr Leu

1 5 10 15

ctt ata ttt ctg gtt gga ata tgg gct gca tgg aaa acc aaa aac agc 96

Leu Ile Phe Leu Val Gly Ile Trp Ala Ala Trp Lys Thr Lys Asn Ser

20 25 30

ggc aac cca gaa gag cac agt gaa gcc atc ata gtc ggg ggc cgt gac 144

Gly Asn Pro Glu Glu His Ser Glu Ala Ile Ile Val Gly Gly Arg Asp

35 40 45

att ggt ttg ttg gtt ggt ggt ttt acc atg aca gcc acc tgg gtt gga 192

Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly

50 55 60

gga ggc tac atc aat ggg aca gca gaa gca gtg tat ggg cca ggt tgt 240

Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Gly Pro Gly Cys

65 70 75 80

ggt cta gct tgg gct cag gca ccc att gga tat tct ctg agt cta att 288

Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile

85 90 95

tta ggt ggt ctg ttt ttt gcg aaa cct atg cgt tcc aag gga tat gtg 336

Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val

100	105	110	
act atg tta gac cca ttt caa cag atc tat gga aag cgc atg ggt ggg			384
Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly			
115	120	125	
ctg ctc ttc atc cct gca ctg atg gga gag atg ttc tgg gct gca gca			432
Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala			
130	135	140	
att ttc tct gca tta ggg gcc acc atc agc gtg atc att gat gtg gat			480
Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp			
145	150	155	160
gtg aac ata tcg gtc att gtc tct gca ctc att gcc att ctt tat acc			528
Val Asn Ile Ser Val Ile Val Ser Ala Leu Ile Ala Ile Leu Tyr Thr			
165	170	175	
cta gtg ggt ggg ctc tac tct gtg gca tat act gat gtt gtc cag cta			576
Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu			
180	185	190	
ttc tgc att ttt ata gga ctg tgg atc agt gtc cct ttt gcc ctg tca			624
Phe Cys Ile Phe Ile Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser			
195	200	205	
cat cct gca gtc acc gac atc gga ttc aca gct gtg cat gct aaa tac			672
His Pro Ala Val Thr Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr			
210	215	220	

cag agt ccc tgg ctg gga acc att gaa tca gtt gaa gtc tac acc tgg 720
 Gln Ser Pro Trp Leu Gly Thr Ile Glu Ser Val Glu Val Tyr Thr Trp
 225 230 235 240

ctt gat aat ttt ctg tta ttg atg ctg ggt gga atc cca tgg caa gcc 768
 Leu Asp Asn Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala
 245 250 255

tac ttc cag agg gtc ctc tct tca tcc tca gcc acc tat gct cag gta 816
 Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val
 260 265 270

ctg tcc ttc ctg gca gct ttt ggg tgc ctg gtc atg gct cta ccc gcc 864
 Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Leu Pro Ala
 275 280 285

ata tgc ata gga gct att gga gct tcc aca gac tgg aac cag act gcc 912
 Ile Cys Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala
 290 295 300

tac ggg tat cca gat ccc aag act aag gag gaa gca gac atg att ctc 960
 Tyr Gly Tyr Pro Asp Pro Lys Thr Lys Glu Glu Ala Asp Met Ile Leu
 305 310 315 320

ccg atc gtt ctg cag tac ctc tgc cct gtg tac atc tcc ttc ttt ggg 1008
 Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly
 325 330 335

ctt ggt gct gtt tca gct gct gtc atg tcc tca gct gac tgc tcc atc 1056
 Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile

340	345	350	
cig tcg gcg agt tct atg ttt gct cgg aat atc tac cag ctt tcc ttc	1104		
Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe			
355	360	365	
aga caa aat gca tca gac aag gaa att gtg tgg gtc atg agg atc act	1152		
Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr			
370	375	380	
gtg ctt gtg ttc gga gca tct gca aca gcc atg gct ttg ctg acg aag	1200		
Val Leu Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys			
385	390	395	400
act gtg tat ggg ctc tgg tac ctg agc tct gac ctt gtc tac atc atc	1248		
Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Ile			
405	410	415	
atc ttc cca cag ctg ctc tgt gta ctc ttc atc aaa gga acc aac act	1296		
Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Ile Lys Gly Thr Asn Thr			
420	425	430	
tat ggg gca gtt gct ggt tat att ttt gga cta ttc ctg aga att act	1344		
Tyr Gly Ala Val Ala Gly Tyr Ile Phe Gly Leu Phe Leu Arg Ile Thr			
435	440	445	
gga gga gag cca tat cta tac ttg cag ccc tta atc ttc tac cct ggt	1392		
Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly			
450	455	460	

tat tac tct gac aag aat ggt ata tac aat cag agg ttc cca ttt aaa 1440
Tyr Tyr Ser Asp Lys Asn Gly Ile Tyr Asn Gln Arg Phe Pro Phe Lys
465 470 475 480

act ctc tcc atg gtt acc tca ttc ttt acc aac att tgt gtt tct tat 1488
Thr Leu Ser Met Val Thr Ser Phe Phe Thr Asn Ile Cys Val Ser Tyr
485 490 495

cta gcc aag tat cta ttt gaa agt gga acc ttg cct cca aaa tta gat 1536
Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp
500 505 510

gta ttt gat gct gtt gtc gca agg cac agt gaa gag aac atg gac aag 1584
Val Phe Asp Ala Val Val Ala Arg His Ser Glu Glu Asn Met Asp Lys
515 520 525

acc att cta gtc aga aat gaa aat atc aaa tta aat gaa ctt gca cct 1632
Thr Ile Leu Val Arg Asn Glu Asn Ile Lys Leu Asn Glu Leu Ala Pro
530 535 540

gtg aaa cct cgg cag agc cta acc ctc agt tca act ttc acc aat aag 1680
Val Lys Pro Arg Gln Ser Leu Thr Leu Ser Ser Thr Phe Thr Asn Lys
545 550 555 560

gag gcc ctc ctt gat gtt gat tcc agt ccg gag ggg tct ggg act gaa 1728
Glu Ala Leu Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu
565 570 575

gat aac tta caa tga 1743
Asp Asn Leu Gln

580

<210> 8

<211> 580

<212> PRT

<213> Mouse

<400> 8

Met Ser Phe His Val Glu Gly Leu Val Ala Ile Ile Leu Phe Tyr Leu

1 5 10 15

Leu Ile Phe Leu Val Gly Ile Trp Ala Ala Trp Lys Thr Lys Asn Ser

20 25 30

Gly Asn Pro Glu Glu His Ser Glu Ala Ile Ile Val Gly Gly Arg Asp

35 40 45

Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly

50 55 60

Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Gly Pro Gly Cys

65 70 75 80

Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile

85 90 95

Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val

100 105 110

Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly
115 120 125

Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala
130 135 140

Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp
145 150 155 160

Val Asn Ile Ser Val Ile Val Ser Ala Leu Ile Ala Ile Leu Tyr Thr
165 170 175

Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu
180 185 190

Phe Cys Ile Phe Ile Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser
195 200 205

His Pro Ala Val Thr Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr
210 215 220

Gln Ser Pro Trp Leu Gly Thr Ile Glu Ser Val Glu Val Tyr Thr Trp
225 230 235 240

Leu Asp Asn Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala
245 250 255

Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val
260 265 270

Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Leu Pro Ala
275 280 285

Ile Cys Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala
290 295 300

Tyr Gly Tyr Pro Asp Pro Lys Thr Lys Glu Glu Ala Asp Met Ile Leu
305 310 315 320

Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly
325 330 335

Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile
340 345 350

Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe
355 360 365

Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr
370 375 380

Val Leu Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys
385 390 395 400

Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Ile
405 410 415

Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Ile Lys Gly Thr Asn Thr
420 425 430

Tyr Gly Ala Val Ala Gly Tyr Ile Phe Gly Leu Phe Leu Arg Ile Thr
435 440 445

Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly
450 455 460

Tyr Tyr Ser Asp Lys Asn Gly Ile Tyr Asn Gln Arg Phe Pro Phe Lys
465 470 475 480

Thr Leu Ser Met Val Thr Ser Phe Phe Thr Asn Ile Cys Val Ser Tyr
485 490 495

Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp
500 505 510

Val Phe Asp Ala Val Val Ala Arg His Ser Glu Glu Asn Met Asp Lys
515 520 525

Thr Ile Leu Val Arg Asn Glu Asn Ile Lys Leu Asn Glu Leu Ala Pro
530 535 540

Val Lys Pro Arg Gln Ser Leu Thr Leu Ser Ser Thr Phe Thr Asn Lys
545 550 555 560

Glu Ala Leu Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu
565 570 575

Asp Asn Leu Gln
580

[Brief Explanation of Drawings]

[Fig. 1]

This is a view showing the result of [^3H] choline uptake of oocytes from *Xenopus* of the present invention being injected with nematode cho-1 (C48D1.3 cRNA) or water.

[Fig. 2]

This is a view showing the result of the effect of Na^+ on choline uptake of oocytes from *Xenopus* of the present invention being injected with nematode cho-1 (C48D1.3 cRNA) or water.

[Fig. 3]

This is a view showing the result of the HC3-induced inhibition of choline uptake of oocytes from *Xenopus* of the present invention being injected with nematode cho-1 (C48D1.3 cRNA) or water.

[Fig. 4]

This is a view showing amino acid sequences of rat CHT1 and nematode CHO-1 of the present invention respectively.

[Fig. 5]

This is a view showing the distribution of neurons expressing cho-1::gfp of the present invention in the nervous system of nematode.

[Fig. 6]

This is a view showing the phylogenetic tree of Na^+ -dependent glucose transporter family.

[Fig. 7]

This is a view showing an expected topology of rat CHT1 of the present invention.

[Fig. 8]

This is a view showing the result of Northern blot analysis of CHT1 mRNA transcript in rat tissue of the present invention.

[Fig. 9]

This is a view showing the result of *in situ* hybridization analysis of CHT1 transcript in a rat brain of the present invention.

[Fig. 10]

This is a view showing the result of *in situ* hybridization analysis of CHT1 transcript in a spinal cord of the present invention.

[Fig. 11]

This is a view showing the result of [^3H] choline uptake of oocytes from *Xenopus* of the present invention being injected CHT1 cRNA of the present invention or water.

[Fig. 12]

This is a view showing the effect of choline concentration on choline uptake in CHT1 of the present invention.

[Fig. 13]

This is a view showing the result of HC3-induced inhibition of choline uptake of CHT1 of the present invention.

[Fig. 14]

This is a view showing the result of Na^+ - and Cl^- -dependent choline uptake of CHT1 of the present invention.

[Fig. 15]

This is a view showing the result of [^3H] HC3 binding to the membrane prepared from COS7 cells being introduced with CHT1 cDNA of the present invention or vector pCDNA 3.1 separately.

[Fig. 16]

This is a view showing the result of saturation analysis of specific [^3H] HC3 binding to the membrane prepared from COS7

cells being introduced with CHT1 cDNA of the present invention or vector pcDNA 3.1 separately.

[Fig. 17]

This is a view showing the result of displacement of specific [^3H] HC3 binding by HC3 of the present invention, choline (Cho), acetylcholine (ACh).


[Name of Document] ABSTRACT

[Abstract]


[The Object] The object of the present invention is to provide a protein having high-affinity choline transporter activity which is important physiologically, a gene encoding the protein, and a method of screening a material promoting the high-affinity choline transporter activity with the use of the same, and the like.

[Solving Means] By examining high-affinity choline uptake activity of Na⁺-dependent transporter cDNA deduced from the genomic sequence of a nematode (*C.elegans*) in a *Xenopus* oocyte expression system, the cDNA (cho-1) of nematode high-affinity choline transporter is identified. Then the cDNA (CHT1) of rat high-affinity choline transporter is cloned from rat spinal cord by using the homology of a base sequence to this cDNA as an index. Similarly, the cDNA of human high-affinity choline transporter is cloned from human genome.


[Selected Drawing] Fig. 4



[Fig. 1]




[Fig. 2]





[Fig. 3]






[Fig. 4]


CHT1	MPFH EGVAT FYLI VGIWAAMKNS----GAEER EALV GGRDIGLLVGGF	56
cht-1	-MAD LGIVAT FYVLI VGIWAGRKSSKELES AGAAEEVILAGRIGTLVGIF	59
I		
CHT1	TMTATWVG GGYINGTAEAY GPGCGLAWAQAP GY SL GGL FAKPMRS GY TMLD	116
cht-1	TMTATWVG GAYINGTAEAY NG GLLGCOAP GY SL GGL FAKMRE GY TMLD	117
II		
CHT1	PFQ IYGKR GGL PALMGEMFWAAAT SALGAT SVI DVDVNISVI SALIA YT	176
cht-1	PFQ KYGQR GGL PALLGETFWTAAI SALGAT SVI G DYNASVT SACIA YT	177
III IV		
CHT1	VGGLY VAYTDVVQLFCIF GLW SVPFA SHPVTDIGFTAVHAKYQSPW GTIES-V	235
cht-1	TGGYY VAYTDVVQLFCIF GLW CVPAAYVHDGAKDISRNAG-----DW GEIGGFK	231
V		
CHT1	EYVTW DNL LLLV GGIPWQAYFQRLSSS SATYAQVLSF AAFGC VMA PAICIGATG	295
cht-1	ETSLW DC LLLV GGIPWQVYFQRLSSKTAHGAQTL SFVAGVGC LMA PPALIGATA	291
VI VII		
CHT1	ASTDWNQTAYGFP PKTKEEAD-----M PLV QYLCPVY SF GLGAVSAAVMSSAD	349
cht-1	RNTDWRM TDYSPWNGTKVES IPPDKRM VPE V QYLTPRV AF GLGAVSAAVMSSAD	351
VIII		
CHT1	SS LSA SMFA NIYQLS RQASD KE IVVMRI V VFGASATAMALLT VYGLWYLS	409
cht-1	SSVLSA SMFA NIKLIRP AS KE IVVMRI V CVGIMATIMALT IOS YGLWYLC	411
IX		
CHT1	DLVY IIFPQLLCVY RKG NTYGA YAGYIFGLFLR TGGEPYQYLQP IFPGYYPDK	469
cht-1	DLVY IIFPQLLCVY MP RS NTYGS LAGYAVGLVLR IGGEPLVSLPA THYPMYT-D	469
X XI		
CHT1	NGIYNQRFPTLSMVIS FTNICVSYLAKYLF SGTLPPLDIE DAVVSR---HSEENM	526
cht-1	G---VQYFPFRITM SSYATIYIVSIOSE KLF SGRLSPWDVYGCVVNIPIDVPLPS	526
XII		
CHT1	DKIQLVRNENIKLNELAPVKPRQS TLSSTF INKEALLDVDSSPEGSGT DNLQ	580
cht-1	DVS FVSS E--TLNMKAPNGTPAP HPNQPS DENTLLHPYSDQSYYSTNSN--	576




[Fig. 5]



[Fig. 6]




[Fig. 7]






[Fig. 8]




[Fig. 9]






[Fig. 10]




[Fig. 11]






[Fig. 12]




[Fig. 13]






[Fig. 14]




[Fig. 15]





[Fig. 16]



[Fig. 17]

